




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

# Journal Pre-proof

Sweet syndrome in post-COVID-19 infection: A case report

Fatima-Zahra Agharbi, Kenza Oqbani, Abir Allaoui, Inas Chikhaoui, Soumiya Chiheb



PII: S1477-8939(21)00229-5

DOI: <https://doi.org/10.1016/j.tmaid.2021.102188>

Reference: TMAID 102188

To appear in: *Travel Medicine and Infectious Disease*

Received Date: 16 October 2021

Revised Date: 18 October 2021

Accepted Date: 19 October 2021

Please cite this article as: Agharbi F-Z, Oqbani K, Allaoui A, Chikhaoui I, Chiheb S, Sweet syndrome in post-COVID-19 infection: A case report, *Travel Medicine and Infectious Disease* (2021), doi: <https://doi.org/10.1016/j.tmaid.2021.102188>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier Ltd.

Letter to the Editor

# **Sweet Syndrome in post-COVID-19 Infection: a Case Report**

Fatima-Zahra Agharbi, Kenza Oqbani, Abir Allaoui, Inas Chikhaoui, Soumiya Chiheb.

Sheikh Khalifa Hospital, Faculty of Medicine, Mohamed VI University of Health Sciences.  
Casablanca, Morocco.

Corresponding author: Agharbi Fatima-Zahra

E-Mail: [aghmarifz@gmail.com](mailto:aghmarifz@gmail.com)

**Keywords:** Sweet syndrome; COVID-19; neutrophilic dermatosis

Dear editor:

The novel coronavirus (SARS-CoV-2), the cause of coronavirus 2019 disease (COVID-19) pandemic, is associated with some cutaneous manifestations. The most frequently reported are: acral lesions, macula-papular rashes, urticarial rashes, vascular lesions and vesicular rashes (1). The occurrence of sweet syndrome in the context of a covid infection of which we report a case is rare; only two cases have been reported in the literature (2, 3). We report here a case of Sweet's syndrome occurring three weeks after a simple coronavirus disease 2019 (COVID-19) infection.

A 61-year-old patient with no particular history who presented three weeks after a simple COVID-19 infection confirmed by positive RT-PCR in nasopharyngeal swabs, made up of fever, cough myalgia and without severe respiratory syndrome, painful erythematous, edematous, purplish plaques in the hands and feet (Figure 1). The biological assessment found an inflammatory syndrome with a CRP of 105 mg, hyperleukocytosis predominantly neutrophilic. The ASLO assay was negative. A thoraco-abdomino-pelvic scanner, performed to rule out associated malignancies, showed no abnormalities. Protein electrophoresis was normal. The patient did not have any history of inflammatory bowel disease or digestive sign, and faecalcal protectin was negative. No supplementary drugs were added to what was indicated above. Pathological examination of a skin biopsy found an inflammatory dermal neutrophilic infiltrate without vasculitis. A Sweet post covid syndrome was retained in view of the clinical and anatomopathological elements. Colchicine treatment was started with marked improvement.

Sweet's syndrome also known as acute febrile neutrophilic dermatosis is a rare neutrophilic dermatosis characterized by a constellation of clinical symptoms, physical features, and pathologic findings which include fever, neutrophilia, tender erythematous skin lesions (papules, nodules, and plaques) most often on face and upper extremities. Histologically, we note a diffuse infiltrate consisting predominantly of mature neutrophils that are typically located in the upper dermis without vasculitis. The cause of this syndrome remains unknown in most cases. However, it has been associated with inflammatory diseases, haematological or visceral malignancies, pregnancy, drug exposure, upper respiratory and gastrointestinal infections (4). Taşkın et al. (2) reported the first case of sweet's syndrome concomitant with SARS-CoV-2 infection in a 61-yearold woman. Simar Berro et al (3) have published sweet's syndrome three weeks after a severe COVID-19 infection. The mechanism of Sweet's syndrome remains unknown; it seems to be related to a hypersensitivity reaction triggered by

infections, neoplastic processes or by drug intake leading to the secretion of cytokines and activation of PNN. Granulocyte colony stimulating factors (G-CSF) is a specific factor in the growth of neutrophils that plays a crucial role in the pathogenesis of Sweet syndrome, however several cases of Sweet syndrome have been reported following administration of GM-CSF (5) and several studies have shown an increase in GM-CSF levels in the acute phase of Sweet syndrome. GM-CSF stimulates the production, activation and chemotaxis of neutrophils that accumulate in the dermis (6, 7). The pathogenesis of sweet syndrome in Covid 19 infection is unclear but is likely the result of cytokine storm and immunologic abnormalities. The aberrant host response in COVID-19 infection was proposed to be centred around neutrophils. In autopsy specimens, neutrophil infiltration in pulmonary capillaries, extravasation of neutrophils to alveolar space and capillaritis with fibrin deposition were shown (8, 9).

We describe here a typical Sweet's syndrome occurring three weeks after COVID-19 infection and we very likely think of a causal link between the two events given that associated malignancy and inflammatory bowel disease were ruled out and the possible involvement of the administered drugs was investigated, but no strong evidence was found in terms of extrinsic and intrinsic imputability.



**Figure 1.** Edematous purplish plaques of hands.

**Acknowledgement**

We are indebted to the patient for giving us the consent for publication.

**Statement of Ethics**

The patient's consent was obtained for the pictures and eventual publication.

**Disclosure Statement**

The authors have no conflicts of interest to declare.

**Funding Sources**

The authors did not receive any funding.

**Author Contributions**

All authors also state that they have read and approved the final version of the manuscript.

**References**

- 1- Recalcati S: Cutaneous manifestations in COVID- 19: A first perspective. J Eur Acad Dermatol Venereol. 2020, 34:212-3
- 2- Berro S, Calas A, Sohier P, Darbord D, Dupin N. Sweet's Syndrome Three Weeks after a Severe COVID-19 Infection: A Case Report. Acta Derm Venereol. 2021; 101 (6): 1-2.
- 3 - Taşkın B, Vural S, Altuğ E, Demirkese C, Kocatürk E, Çelebiİ, et al. COVID 19 Presenting with Atypical Sweet's Syndrome. J Eur Acad Dermatol Venereol. 2020; 10 : 534-35. PMID: 32452542
- 4 - Caroline Lang , Sandrine Quenan . Sweet's syndrome, a diagnosis to keep in mind. Rev Med Suisse, 2017 Mar 29;13(556):678-683.
- 5- KR. Arbetter , KW.Hubbard , SN. Markovic , LE.Gibson, RL. Phyliky. Case of granulocyte colony-stimulating factor-induced Sweet's syndrome. Am J Hematol. 1999;61(2):126-9.
- 6- MA. Reuss-Borst, CA. Muller , HD.Waller. The possible role of G-CSF in the pathogenesis of Sweet's syndrome. Leuk Lymphoma. 1994;15(3-4):261-4.
- 7 – MA. Reuss-Borst, G. Pawelec , JG. Saal, HP. Horny , CA. Muller , Waller HD. Sweet's syndrome associated with myelodysplasia: possible role of cytokines in the pathogenesis of the disease. Br J Haematol. 1993; 84(2):356-8.
- 8- Barnes BJ, Adrover JM, Baxter-Stoltzfus A et al. Targeting potential drivers of COVID-19: Neutrophil extracellular traps. J Exp Med 2020; 217.
- 9- Fox SE, Akmatbekov A, Harbert JL, Li G, Brown JQ, Vander Heide RS. Pulmonary and cardiac pathology in Covid-19: the first autopsy series from New Orleans, 2020.